REFERENCES/ BIBLIOGRAPHY

1. Andersen ML, Santos HER, Seabra MV, Silva ABS, Tufik S.,2004.Evaluation of acute and chronic treatment with Harpagophytum procumbens on Freund's adjuvant-induced arthritis in rat.Journal of Ethnopharmacology 91, 325-330.
2. Ali BH, Blunden G. Pharmacological and toxicological properties of Nigella sativa Phytother Res .2003 ; 17 (4): 299-305.
3. Akbar Hajizadeh Moghaddam, Seyed And Mohammad Ali brahimzadeh. Antioxidant, antihemolyticand nephroprotective activity of aqueous extract of Diospyros lotus seeds. actapoloniae pharmaceutica n drug research,2012.69(4),687-692.
4. Al-Jassir MF. Chemical composition of microflora of black cumin (Nigella sativa L.) seeds growing in Saudi Arabia. Food Chem . 1992 ; 45 : 239-242.
5. Abdel-Fattah M, Matsumoto K, Watanabe H. Antinociceptive effects of Nigella sativa oil and its major components in mice. Eur J Pharmacol . 2000 ; 400 : 89-97.
6. Andersen M.L., D'Almeida V., Ko G.M., Martins P.J.F., Tufik S. (2016) Care and Maintenance of Laboratory Animals. In: Andersen M., Tufik S. (eds) Rodent Model as Tools in Ethical Biomedical Research. Springer, Cham.
7. A.M. Bendele: Animal models of rheumatoid arthritis. J Musculoskel Neuron Interact 2001; 1(4):377-385.
8. Badary OA, Taha RA, Gamal El-Din AM, Abdel-Wahab MH. Thymoquinone is a potent superoxide anion scavenger. Drug Chem Toxicol . 2003 ; 26 (2): 87-98.
9. Burits M, Bucar F. Antioxidant activity of Nigella sativa essential oil. Phytother Res . 2000 ; 14 (5): 323-328.
10. Creasy DM. (2002) Histopathology of the male reproductive system I I: interpretation. Curr Protoc Toxicol. Nov;Chapter 16:Unit16.4.
11. Devi Swapna P V 2005 Practical Pharmaceutical Chemistry, Part 24th Edition Cbs Publications And Distributors, 2005, ISSN 157-168.(book).

12. Gad,S.C. (1982). A Neuromuscular Screen for Use in Industrial Toxicology.J.Toxicol Environ.Health,9,691-704.
13. El-Kadi A, et al. 1st International Conference on Scientific Miracles of Quran and Sunnah, Islamabad, Pakistan.
14. El Tahir KE, Ashour MM, al-Harbi MM. The respiratory effects of the volatile oil of the black seed (Nigella sativa) in guinea-pigs: elucidation of the mechanism(s) of action. Gen Pharmacol . 1993 ; 24 (5): 1115-1122.
15. Effect of Chlorophytum borivilianum on adjuvant induced arthritis in rats. Annals of Biological Research, 2010, 1 (1) : 36-40.
16. Firestein G.S. (2003). Evolving concepts of rheumatoid arthritis. Nature., 423: 356–361.
17. Ghazanfar SA. Handbook of Arabian Medicinal Plants . Boca Raton, FL: CRC Press; 1994:180-181.
18. Ghannadi A, Hajhashemi V, Jafarabadi H. An investigation of the analgesic and anti-inflammatoty effects of Nigella sativa seed polyphenols. J Med Food . 2005 ; 8 (4): 488-493 .
19. Hajhashemi V, Ghannadi A, Jafarabadi H. Black cumin seed essential oil, as a potent analgesic and anti-inflammatory drug. Phytother Res . 2004 ; 18 (5): 195–199.
20. Hong Li, Yun-Yi Zhang*, Hui-Wen Tan, Yong-Feng Jia, Duan Li: Therapeutic effect of tripterine on adjuvant arthritis in rats. Journal of Ethnopharmacology 118 (2008) 479– 484.
21. Halliwell B. (1995). Oxygen radicals, nitric oxide andhuman inflammatory joint disease. Annals of RheumaticDisease., 54: 505–510.
22. H.B. Townsend, K.G. Saag: Glucocorticoid use in rheumatoid arthritis:Benefits, mechanisms, and risks.
23. Ibrahim Tekeoglu1*, Ali Dogan2, Levent Ediz1, Mustafa Budancamanak1and Adnan Demirel1: Effects of Thymoquinone (Volatile Oil of Black Cumin) on Rheumatoid Arthritis in Rat Models. Phytother. Res. 21, 895–897 (2007).

24. Keshri G, Singh MM, Lakshmi V, Kamboj VP. Post-coital contraceptive efficacy of the seeds of Nigella sativa in rats. Indian J Physiol Pharmacol . 1995 ; 39 (1): 59-62.
25. Khohinur Hossain 2013 A Validated RP-HPLC Method for Simultaneous Estimation of Antidiabetic Drugs Pioglitazone HCl and Glimepiride angladesh Pharmaceutical Journal 16(1): 69-75, 2013.
26. International Conference On Harmonization**:** Ich Q 2 (R1) Validation Of Analytical Procedures: Text And Methodology 1995.(book).
27. International council for harmonization (ICH) harmonized tripartite guidelines M3 (R2), guidance on nonclinical safety studies for the conduct for human clinical trials and marketing authorization for pharmaceuticals, adopted on June 2009.
28. International council for harmonization (ICH) harmonized tripartite guidelines S3A, note for guidance on toxickinetics: the assessment of systemic exposure in toxicity studies, adopted on 27 October 1994 by the ICH steering committee.
29. Linos AA, Worthington JW, Ofallon WM and Kurland LT: The epidemiology of rheumatoid arthritis in Rochester, Minnesota: a study of incidence, prevalence andmortality. Am J Epidemiol 1980; 111:87–98.
30. Lee DM and Weinblatt ME: Rheumatoid arthritis. The Lancet 2001; 358: 903–911.
31. L.N. Babu and Manoj G. Tyagi: Evaluation Of The Anti-Inflammatory Effect of Dipyridamole, Atp *Per Se* And In Combination With Methotrexate in Freund's Adjuvant Induced Arthritis In Rats. Recent Research in Science and Technology 2010, 2(3): 71–75.
32. M. Rassol and E.P. Sabina: Antiinflammatory Effect of the Indian Ayurvedic Herbal Formulation Triphala on Adjuvant-induced Arthritis in Rats. Phytother. Res. 21, 889–894 (2007).
33. Mukher je, P.K., Shau, M., Suresh, B.,Evaluation of Indian traditional medicine. Eastern Pharmacist ,1998, 42(8) 21-24.

34. Mangesh s Bansod *,1, Virendra g. Kagathara 1, 2, Rohini r. Pujari 1, 3, vivek b. Patel 2, hardik h.ardeshna 2: therapeutic effect of a polyherbal preparation on adjuvant induced arthritis in wistar rats. International journal of pharmacy and pharmaceutical sciencevol 3, suppl 2, 2011.
35. Nickavar B, Mojab F, Javidnia K, Amoli MA. Chemical composition of the fixed and volatile oils of Nigella sativa L. from Iran. Z Naturforsch . 2003 ; 58 (9-10): 629-631.
36. OECD, 1995: OECD N° 407, "Repeated Dose 28-day Oral Toxicity Study in Rodents". The Organisation for Economic Co-operation and Development (OECD) Guidelines for the Testing of Chemicals, Adopted by the council on 27th July, 1995.
37. OECD, 2001: OECD N° 425, "Acute Oral Toxicity-Up-and-Down Procedure". The Organisation for Economic Co-operation and Development (OECD) Guidelines for the Testing of Chemicals, Adopted by the Council on 17th December, 2001.
38. OECD (2006). Report of the Validation of the Updated Test Guideline 407 Repeat Dose 28-day Oral Toxicity Study in Laboratory Rats. Environmental Health and Safety Publications Series on Testing and Assessment No. 59. Organisation for Economic Cooperation and Development, Paris.
39. OECD (2000) Guidance document on the recognition, assessment and use of clinical signs as humane endpoints for experimental animals used in safety evaluation. Series on Testing and Assessment No 19. ENV/JM/MONO(2000)7.
40. Perera Pathirage-Kamal1, PENG Cheng1, LV Xue 1, LI Yun-Man1*,FANG Wei-Rong1, HAN Cai-Feng2: Effects of *Yishen Juanbi* (YJB) Pill on ExperimentalRheumatoid Arthritis. *Chinese Journal of Natural Medicines* 2010, 8(1): 0057−006.
41. Salem ML. Immunomodulatory and therapeutic properties of the Nigella sativa L. seed. Int Immunopharmacol . 2005 ; 5 (13-14): 1749-1770 .
42. Siavash Parvardeh and Mohammad Fatehi: Effects of Thymoquinone, the Major Constituent of Nigella sativa Seeds, on the Contractile Responses of Rat vas Deferens. Pharmaceutical Biology 2003, Vol. 41, No. 8, pp. 616–621.

43. Spadoni G, Bedini A, Rivara S, Mor M , 2011 Melatonin receptor agonists: new options for insomnia and depression treatment , CNS Neurosci Ther. 2011 Dec;17(6):733-41.

44. Suha Abudoleh, 1Ahmad Disi, 2Eyad Qunaibi and 3Talal Aburjai: Anti-Arthritic Activity of the Methanolic Leaf Extract of *Urtica pilulifera* L. on Albino Rats. American Journal of Pharmacology and Toxicology 6 (1): 27-32, 2.

45. Sonali P Desai* and Daniel H Solomon: The use of glucocorticoids in rheumatoid arthritis -no 'rational' approach yet. Desai and Solomon *Arthritis Research & Therapy* 2010, **12**:127.

46. Weingand K, Brown G,Hall Retal. (1996). "Harmonisation of animal Clinical Pathology Testing in Toxicity and Safety Studies", Fundam. & Appl. Toxicol., 29:198-201.